Steven M. Ruben
Appl. No. 10/662,429



BEST AVAILABLE COPY

Ruben EXHIBIT #61

Department Molecular Biology

Subject

Name Guo-Liang Yu II

Address

National Brand 43-648

Computation Notebook

Dennison Stationery Products Co., Framingham, MA 01701

0 73333 43648 8

75 Sheets
11¾" x 9¼"
4x4 Quad.

Ruben EXHIBIT 2061
Ruben v. Wiley et al.
Interference No. 105,077
RX 2061

8/21/95 Two New TNF: HPDD012 TNF Epsilon

HETBT71 TNF delta maybe splice variant

there are differences between the two clones

Oligo captrue cloned full length Named HETBT71SO7

design oligos for seq Rpo2 CTCCCAGCTTGGAAGACCAG

Rpo3 TTCTGCATGCCCACCTCTC

Fpo6 GAACAGAAGCAAGATATCCG

design oligos for δ-specific Cintron J?

design oligos for construct into pQE70

TNFδ-SphI: CGCGCATGCAGGGGACAGGAGCC

TNFδ-HindIII: CGCAAGCTTACAATCACAGTTTCACAAC

TNFδ-specific oligo is given to L. Xing for Northern

8/25/95 PCR to generate insert DNA for following constructs

1. TNF5 SphI + HindIII ⟹ pCC70
2. TNFe SphI + HindIII ⟹ pCC70
3. TNFr dig NcoI + HindIII ⟹ pCC60
4. TNFr FL BamHI + Bam H3' ⟹ N346
5. TNFr FL BamH + Bam H3' ⟹ New C40 vector

10 λ 10X PCR buf BM
10 λ 2mM dNTP
oligos
0.5 μl PWO polymerase
100 μl

95C 2' 30[95C 1' 55C 1' 72C 1'] 72C 5 min

PCR ⟹ low melting gel (Nusieve)

± 4 PCR for C40 clone did not work

repeat PCR

SUPERVISOR- Patrick J. Stiller
DATE- 08/31/95

9/1/95 PCR ~~to~~ product BamHI fragment for TNFR

& extract 2X

cells est

ethanol ppt

I cut -- check 1Δ on gel

give to Lily for clone into

CHO vectors

9/6/95

Summary on subtraction using biotinylated drivers generated by PCR using biotin-dCTP

	Driver	PRL	G4	mito
HPTA	—	6.5%	27%	—
HPTG	G4 PL	1.6%	18%	—
HPTM	G4 PL	1.2%	15%	—
HSDB	—			34%
HSDS	mito			28%
HKIA	—			45%
HKIS	mito			23%


Subtraction result
% In ESTs
30
20
10
0
HPTA
HPTG
HPTM
growth hormone
prolactin


Subtraction
% In EST
50
40
30
20
10
0
HKIA
HKIS
mitochondrial

9/6/95 Modify the approach:

① using regulator drop to generate ss DNA

② photo biotin DNA
this is proven to be efficient

③ Try both Long hybridization at 42°C o/n
and short hybridization at RT 1 hr

④ remove biotin by extract add one more step should
no difference by using magnetic beads

9/15 Discussion w/ Fouad — idea



NORTHERN BLOT DATA SHEET

OPERATOR: Lily Xing DATE 9/1/94

LANE #	NAME	RNA (µl/10µg)	Note
1	Brain	20 ug	
2	kidney	"	
3	small intestin	"	
4	testis	"	
5	pancreas	"	
6	prostate	"	
7	Heart	"	
8	Liver	"	
9	Lung	"	
10	thymus	"	
11	spleen	"	
12	placenta	"	
13	Colon	"	
14	ovary	"	
15	Leukocyte	"	
16	mucle	"	
17			
18			
19			
20			

NOTE:

#3 gel



NORTHERN BLOT DATA SHEET

OPERATOR: Lily Xing DATE 9/1/94

LANE #	NAME	RNA (µl/10µg)	Note
1	Brain	20 ug	
2	kidney	"	
3	small intestin	"	
4	testis	"	
5	pancreas	"	
6	prostate	"	
7	Heart	"	
8	Liver	"	
9	Lung	"	
10	thymus	"	
11	spleen	"	
12	placenta	"	
13	Colon	"	
14	ovary	"	
15	Leukocyte	"	
16	muscle	"	
17			
18			
19			
20			

NOTE:

#3 gel



9/18/95 Design oligos for SOD4 to determine which ATG is the real ATG

3 oligos AAATTAACCCTCACTAAAGGGCCATCATGGCAGCGGCCGG

T3 - SOD ATG3

SODATG2 ATG GTCTTGGTGCAC

SODATG1 ATGACCTGTCAGAGC

PCR to generate pcr product for TNT

3' oligo SODEND

CGTCTAGAGGTCCTGCTCAAAGGTGGG





SOD4 T3 + T7 PCR

9/19 Subtraction Result

Sequence of two New HPT Subtracted library Come back

in HPTY which is long hyb (O/N at 42°C)

GH reduced to 10% prolactin to 0%

HPTX — short hyb is not as good

GH → 15% prolactin 3%

→ try hybridization with 5% PEG

9/20/95 TNT promega kit

DNA: pBluescpt sk⁻
SOD full length in pBluescpt
PCR product from ΔTG1
ΔTG2
ΔTG3

T7 - CSG16 in pA2

make a premix
- 12.5 ml Rabbit Reticulocyte lysate
- 0.1 ml bf
- 0.5 ml T3 or T7 RNA polymerase
- 0.5 ml Amino Acid - met
- 2 ml 35S-met
- 0.5 λ RNase inhibitor from Boehringer
- 3 λ DNA
- 8 λ ddH2O

30°C 1hr



9/25 repeat TNT

10 λ reaction

2 nl PCR product similar amount

Full length SOD1	— 0	?
ΔTG1	— 1	27 Kd
ΔTG2	— 2	24 kd
ΔTG3	— 3	21 Kd

30°C 1 hr

add 10λ loading dye boil for 2'

Spin Load 10λ on 10% SDS gel



This experiment indicate there is another ATG upstream maybe the real ATG

Check sequence there is another 19 AA upstream this will make the protein 29 K

SUPERVISOR

DATE- 09/27/95

10/4/95

Subtraction result come back with a nice improvement by add 5% PEG (8000) in hybridization mix. it may be increase hybridization efficiency

result:		GH	pr
	before subtraction :	2%	7%
	after subtraction :	5%	1%

— clone Full length TMFepsilon by PCR Hpp 1. brain

Hpp was mass excised and used to PCR using

TM13 Reverse + HLTBT71Fp06
+ HLTBT71Fp04

one band of 0.8 Kb seen

— Second PCR

1λ first PCR T3 + HLTBT71Fp05
Fp04

— PCR to construct SUMO expression vectors





- Lily will clone into pQE60

10/5/95 Gene trapper

genes: TNFR p55
TNF5
TGF β — WWH
Rad 16 — YFW

Label oligos: w/ TDT

TGF β 11362
11356

TNF-5 TNF5 Rp01

TNFR p55 cap1
+
Rp-2

Rad16 - R8
+
R11

Cabling nd working



produce SS DNA by gene T cDNA

Clontech library



TNF5 → Leukocyte

TGFβ → M

Rad 16 → L

TNFR p55 →

16/6/95 Repeat biotin labeling of oligos

ethanol ppt oligos final concentration 200ng/l

3μg DNA (12 μl)
10 λ 5x TdT bfr
5 λ biotin-ddUTP
8 λ TdT



working

10/9/95 Capture

add 6 λ 4x bfr — 95°C 1'
add 5 λ TNFR oligo + Brain
0.5 λ other oligos + Leukocyte — TNFS
Leukocyte — Rad6
Brain — TGFβ

Hyb:

37°C 1 hr

treat SA - Magnetic beads, capture, wash

Repair all 10 λ

10/9/95

transfor to DH10B from BRL

plate	10λ	100λ
TNFRp55	—	40 colony
TNF5	—	41
TGFβ	—	1
Rad 1b	—	6

add 1ml 2x Freeze liqu to remaining

10/10/95 PCR to identify positive clones.

2 gene specific primers

TNF5

50% clones are positive

2 gene specific primers

TNFRp55



~5% clone positive

not so colony hybr

Show only 32/100

hybridize

Sap + TNF5 Fp1



10/16/95 Fusion construct for TNFr with IL6 signal

PCR:

Oligos: IL6 BamH1

CGC GGA TCC ATG AAC CAT GAA CTC CTT CTC CAC

Met

IL6-TNFr1 IL6 TNFr

IL6-TNFr2



50 µl reaction 58 program

Fusion for Full length TNFr (HPDD012)

PCR from HpD library



SUPERVISOR- [signature]

DATE- 10/17/95

10/23 Seq analysis of 4 HTTBN61 clones

HTTBN61 S08 — wrong clone
HTTBN61 S23 — wrong clone
HTTBN61 S02 — wrong clone
HTTBN60 S07 — wrong clone

design new primers for oligo capture ?

— have Lily do a colony hybridization

— the construct of IL6/TNFr is missing kozak sequence

new oligo is made — IL6's'

GCG GG ATCC GCCACC ATG TCC TTC TCC AC
BamHI kozak MET

PCR to generate — IL6/TNFr fusion.

Clones are made into the two CHO vectors by Lily

10/25 still having problem get HPDD12 PCR fragment

PCR products





Low melting gel
Φ 2x
$CHCl_3$

digest w/ Hind III + BamH I

Clone it pbluescript Hind III + BamH I

10/26

PCR products 5μl/50



	primer	template
1	T3 + F2	HPDD12
2	T7 + R3	HPDD12
3	TNFδ H3 + R3	HPDD12
4	T3 + F2	— PCR① — HPD mutant
5	T3 + F2	HPD
6	T7 + R3	HPD
7	TNFδ H3 + R3	HPD
8	T3 + F2	— PCR② — HPD mutant

135

10/17 Low melting gel



Cut out DNA

PCR 4 → 0.6 - 1.1 Kb

PCR 8 - -

R3+Hd3 →
(PCR 7)

Φ 2x

Ligase 1x

ethanol ppt

mix PCR 4 + PCR 7
PCR 8 + PCR 7

PCR using T3 + Hd3



← low melting gel purify top band

HindIII + BamHI digestion

Clone into pBluescript HindIII + BamHI digestion

PCR check → no clone found

11/7/95

oligo flanking TNFδ Fpo7 + Rpo7

PCR	DNA	primer
①	HPD l. brary	Rpo7 + Fpo7
②	HPD l. brary	Rpo7 + Fpo7 + TNFδ Bam + TNFδ Hind III
③	HPD l. brary	TNFδ BamHI + Hind III
④	HLTBT71 sq	Rpo7 + Fpo7 expect → 1.7 kb
⑤	HLTBT71 sq	5 Bam + 3 Hind III expect 600 bp

1 2 3 4 5



wrong size for 3 Bam product

11/10 New primer made for TNFδ BamHI

PCR	DNA	primer
1)	HPD l. brary	Rpo7 + Fpo7
2)	HPD l. brary	5 BamHI + Hind III
3)	PCR ①	Rpo7 + Fpo7
④	PCR ①	BamHI + Hind III
5)	HLTBT71 sq	Rpo7 + Fpo7
6)	HLTBT71 sq	5 BamHI + Hind III

1 2 3 4 5 6



11/14/95 2% low melting gel purification PCR 2 & PCR C

D/ QIAEX extr ethanol ppt

digest with BamHI + HindIII

also digest: Vector pQE-9

Ligation o/n

Design oligos to test ligation for SAGE

Ligation oligo 1 GAGTCAGTTCAATGCCAAACGGCATG

Ligation oligo 2 CCGTTTGGCATTGAACTGACTCCATC

SUPERVISOR- Patrick O. Brown

DATE- 11-20-95

11/20/95 RACE for HTTBN61 TNFR p55 homolog

The following library has the gene based on PCR using 2 gene specific primers

HCQ HLQ HFV HFB HPD HAP HTX HE9
1 2 3 4 5 6 7 8

MisRv T3

F3 F4

PCR using MisRv + F4

HPD
9 = TNFeRpo8 + Fpo7
10 : Rpo8 + Fpo6
11 : Rpo8 + δHindIII



Second PCR

T3 + F3

1 – 8 template PCR 1-8

9 HTTBN61

10 HTTBN61 T7 + Rpo1

11/21/95



5λ + 10% (gel purified)

11/28

T3 + T7 PCR



No specific products

11/28 TMS His tag construct

Seq verified digestion w/ BamH HindIII



Signed by RG and SR

gave DNA to protein engineers



12/6/95 prepare linkers for SAGE

large scale oligo synthesis each 7.5 mg

linker A

TTTTAATTAACCCTCACTAAAGGGCTCGCACGGATGCATG 4368
TTAATTGGGAGTGATTTCCCGAGCGTGCCTAC 4370

linker B

TTTTGTAATACGACTCACTATAGGGCAAGTCGGATGCATG 4369
CATTATGCTGAGTGATATCCCGTTCAGCCTAC 4371

— 12% urea acrylamide gel purify the oligos
put through C18 column

— or ϕ/CHCl3 ext ethanol ppt + equal V — 1M Tris pH 7.5
3 V 60% EtOH

Concentration

Sample ID	abs 260.0 nm	abs 280.0 nm		260.0 nm / 280.0 nm	260.0 nm / 280.0 nm	
4368	0.3014	0.1908	4.5 ml	1.5797	0.6330	ϕ extracted
2 69	0.3548	0.2081	5.3	1.7045	0.5867	
3 70	0.3056	0.1975	4.5	1.5468	0.6465	
4 71	0.3139	0.2102	4.8	1.5171	0.6591	
5 68	0.0762	0.0489	1.1	1.5581	0.6414	gel purified
6 69	0.0768	0.0456	1.2	1.7304	0.5778	
7 70	0.1143	0.0757	1.7	1.5114	0.6616	
8 71	0.1009	0.0660	1.5	1.5292	0.6539	
9						

— Anneal

80 μg (4368) + 64 μg (4370) + 20 λ 10x Kinase buffer
80 μg (4369) + 64 μg (4371) — 70 mM Tris Cl
10 mM MgCl
1 mM DTT

add H2O to 200 λ

A B gel purified

A' B' ϕ/cuts/ethanol ppt

65°C → cool to 42°C put heat block on bench

add 1λ 100 mM ATP
1λ 100 mM ddATP
2λ PNK
2λ Klenow

37°C 1 hr

take 2λ each ligate in 10λ 2λ 5× BRL ligation buf
1λ NEB high conc ligase

RT 2-3 hrs

12% native acrylamide gel



1	4368	ϕ/cuts ethanol ppt
2	4370	
3	4369	
4	4371	
5	4368	gel purified
6	4370	
7	4369	
8	4371	
9	LinkerA	self ligation
10	LinkerA'	
11	B	
12	B'	
13	LinkerA	
14	A'	
15	B	
16	B'	

12/12/95 Northern fox HTTB#64 TMTR

1 2 3 4 5 6 7 8 9 10 11 12 13
Eye Heart Liver Lung Colon Stomach Pancreas Kidney Intestine Testicle Thymus Gallbladder
14. Kidney





12/11/95 CAGE

Linkers A and linkers B

13
Gallbladder



HBC library plasmid DNA digest w/ XhoI
total 7 µg



take ½ (3 µg)
↓ 90°C 5 min
add 0.5 µg biotinylated oligo dT LINKER
↓ 37°C
Klenow 1X 100 mM dNTP
30 min



↓ digest w/ NlaIII



↓ ligation in 20A PKL buffer
RT 3 hr

Linker A
0.2 µg

Linker B
1.2 µg

12/12 pre-wash ... streptavidin magnetic beads

TE wash 3X

Resuspend in 30 λ TE. add ligation.

bind 1 hr. at RT mix occasionally

wash w/ TE/1m NaCl 2X. transfer to new tube

wash 2X more

digestion with FokI at 37°C.



Φ/cuts elute pH↓ T4 DNA poly inverse

+

1λ 100mM dNTP

RT 30'

↓ ligation

8°C

mix A + B o/n

↓ PCR T3 + T7 66 program

0.1 1. 4λ ligation as template (30 cycles)



12/12 prewash strepavidin magenetic beads

TE wash 3X

resuspend in 30λ TE add ligation

Bind 1 hr at RT mix occasionaly

wash w/ TE/1M NaCl 2X, transfer to new tube

wash 2X more

digestion with FokI at 37°C



Φ/CCl3 ethanol ppt ↓ T4 DNA polymerase + 1λ 100mM dATP

RT 30'

↓ Ligation 8°C o/n

mix A + B

↓ PCR T3 + T7 66 program (30 cycles)

0.1 1. 4λ ligation as template

0.1 1. 4



5c





This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☑ FADED TEXT OR DRAWING
- ☑ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☑ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☑ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.
As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.